

Routine Childhood Immunisation Programme

| When to immunise | Diseases vaccine protects against | How it is given |
|---|---|---------------------------------------|
| 2 months old | Diphtheria, tetanus, pertussis (whooping cough), polio and Hib | One Injection |
| | Pneumococcal Infection | One Injection |
| | Rotavirus | Orally |
| | Meningococcal B Infection | One Injection |
| 3 months old | Diphtheria, tetanus, pertussis, polio and Hib | One Injection |
| | Meningococcal C Infection | One Injection |
| | Rotavirus | Orally |
| 4 months old | Diphtheria, tetanus, pertussis, polio and Hib | One Injection |
| | Pneumococcal Infection | One Injection |
| | Meningococcal B Infection | One Injection |
| Just after first birthday | Measles, Mumps and Rubella | One Injection |
| | Pneumococcal Infection | One Injection |
| | Hib and meningococcal C Infection | One Injection |
| | Meningococcal B Infection | One Injection |
| Every year from 2 years old up to P7 | Influenza | Nasal spray or injection |
| 3 years and 4 months old | Diphtheria, tetanus, pertussis and polio | One Injection |
| | Measles, Mumps and Rubella | One Injection |
| Girls 12 to 13 years old | Cervical Cancer caused by human papillomavirus types 16 and 18 and genital warts caused by types 6 and 11 | Two or three injections over 6 months |
| 14 to 18 years old | Tetanus, Diphtheria and Polio | One Injection |
| | Meningococcal ACWY | One Injection |

